

On the Bourque–Ligh Conjecture of Least Common Multiple Matrices

Shaofang Hong

metadata, citation and similar papers at core.ac.uk

Peoples Republic of China

Communicated by Walter Feit

Received October 2, 1996

Let $S = \{x_1, \dots, x_n\}$ be a set of n distinct positive integers. The matrix $[S]_n$ having the least common multiple $[x_i, x_j]$ of x_i and x_j as its i, j -entry is called the least common multiple (LCM) matrix on S . A set S is gcd-closed if $(x_i, x_j) \in S$ for $1 \leq i, j \leq n$. Bourque and Ligh conjectured that the LCM matrix $[S]_n$, defined on a gcd-closed set S , is nonsingular. In this paper we prove that the conjecture is true for $n \leq 7$ and is not true for $n \geq 8$. So the conjecture is solved completely.

© 1999 Academic Press

I. INTRODUCTION

Smith [8] showed that the determinant of the $n \times n$ matrix $[(i, j)]_n$ which has the greatest common divisor (i, j) of i and j as its i, j -entry is the product $\phi(1)\phi(2) \cdots \phi(n)$, where ϕ is Euler's totient function. He also proved that if f is an arithmetical function and $[f(i, j)]_n$ is the $n \times n$ matrix having f evaluated at the greatest common divisor (i, j) of i and j as its i, j -entry, then $\det[f(i, j)]_n = (f * \mu)(1)(f * \mu)(2) \cdots (f * \mu)(n)$, where μ is the Möbius function and $f * \mu$ is the Dirichlet convolution of f and μ . In 1972, Apostol [1] extended Smith's result. In 1988, McCarthy [7] generalized Smith's and Apostol's results to the class of even functions (mod r). In 1993, Bourque and Ligh [4] extended the results of Smith, of Apostol, and of McCarthy.

Let $S = \{x_1, \dots, x_n\}$ be a set of n distinct positive integers. The matrix $(S)_n$, having the greatest common divisor (x_i, x_j) of x_i and x_j as its i, j -entry, is called the greatest common divisor (GCD) matrix on S . The matrix $[S]_n$ having the least common multiple $[x_i, x_j]$ of x_i and x_j as its i, j -entry is called the least common multiple (LCM) matrix on S . The set



S is called factor-closed if it contains every divisor of x for any $x \in S$. Clearly, for any positive integer n , the set $\{1, 2, \dots, n\}$ is factor-closed. A set S is called gcd-closed, if $(x_i, x_j) \in S$ for $1 \leq i, j \leq n$. Obviously, a factor-closed set is gcd-closed but not conversely. For example, $S = \{1, 2, 3, 8, 10\}$ is gcd-closed but not factor-closed.

In 1989, Beslin and Ligh [2] investigated the structure of the GCD matrix $(S)_n$ on a set S of distinct positive integers. In 1992, Bourque and Ligh [3] obtained formulas for the inverses of the GCD matrix $(S)_n$ and the LCM matrix $[S]_n$ on a factor-closed set S . In the meantime they obtained a formula for the inverses of the GCD matrix $(S)_n$ and a formula for the determinant of the LCM matrix $[S]_n$ on a gcd-closed set as

$$\det[S]_n = \prod_{k=1}^n x_k^2 \alpha_k, \quad (1)$$

where

$$\alpha_k = \alpha_k(x_1, \dots, x_k) = \sum_{\substack{d|x_k, \\ d \nmid x_i, \\ x_i < x_k.}} g(d), \quad (2)$$

with the arithmetical function g defined by $g(m) = \frac{1}{m} \cdot \sum_{d|m} d \cdot \mu(d)$, and the function μ is the Möbius function. In [3], Bourque and Ligh raised the following

Conjecture. The LCM matrix $[S]_n$ on a gcd-closed set S is nonsingular.

Recently, Haukkanen, Wang, and Sillanpää [5] gave a counterexample for $n = 9$. In [6] we show that the conjecture is true for $n \leq 5$.

In this paper we introduce a new method to reduce, greatly, the formula for α_k . We prove that the conjecture is true for $n \leq 7$ and is not true for $n \geq 8$. So the conjecture is solved completely.

II. THE MAIN LEMMAS

Throughout this section, let the set $S = \{x_1, \dots, x_n\}$ of n distinct positive integers be gcd-closed and $1 \leq x_1 < \dots < x_n$. Let $|A|$ denote the cardinality of any finite set A .

LEMMA 1 ([6]). For $1 \leq k \leq n$, we have

$$\alpha_k = \beta_k(x_1, \dots, x_k), \quad (3)$$

where the function β_k on a set $Z = \{z_1, \dots, z_k\}$ of k distinct positive integers is defined as

$$\beta_k(z_1, \dots, z_k) = \frac{1}{z_k} + \sum_{r=1}^{k-1} (-1)^r \cdot \sum_{1 \leq i_1 < \dots < i_r \leq k-1} \frac{1}{(z_{i_1}, \dots, z_{i_r}, z_k)}, \quad (4)$$

where $(z_{i_1}, \dots, z_{i_r}, z_k)$ denotes the greatest common divisor of $z_{i_1}, \dots, z_{i_r}, z_k$.

Therefore we can compute α_k by either (2) or (3). It is obvious that formula (3) is easier to use than (2). Now we give further reduction for the formula of α_k .

LEMMA 2. Let $T_k = \{x \in S \mid x < x_k \text{ and } x \nmid x_k\}$, $1 \leq k \leq n$. Let $\{x_1, \dots, x_k\} \setminus T_k = \{x_{k,1}, x_{k,2}, \dots, x_{k,r_k}\}$, where $x_1 = x_{k,1} < x_{k,2} < \dots < x_{k,r_k} = x_k$, $2 \leq r_k \leq k$. Then

$$\alpha_k = \beta_{r_k}(x_{k,1}, x_{k,2}, \dots, x_{k,r_k}). \quad (5)$$

Proof. If $T_k = \phi$, then it follows from Lemma 1 that (5) holds. If $T_k \neq \phi$. Let $x_t \in T_k$. Then $x_t < x_k$. Let $\{x_1, \dots, x_k\} \setminus \{x_t\} = \{x'_1, \dots, x'_{k-2}, x_k\}$, where $x'_1 < \dots < x'_{k-2} < x_k$. Since

$$\beta_k(x_1, \dots, x_k) = \frac{1}{x_k} + \Delta'_t + \Delta_t, \quad (6)$$

where

$$\Delta'_t = \sum_{r=1}^{k-2} (-1)^r \cdot \sum_{\substack{1 \leq i_1 < \dots < i_r \leq k-1 \\ i_j \neq t}} \frac{1}{(x_{i_1}, \dots, x_{i_r}, x_k)}, \quad (7)$$

$$\Delta_t = \sum_{r=0}^{k-2} (-1)^{r+1} \cdot \sum_{\substack{1 \leq i_1 < \dots < i_r \leq k-1 \\ i_j \neq t}} \frac{1}{(x_{i_1}, \dots, x_{i_r}, x_t, x_k)}, \quad (8)$$

by the definition of β_{k-1} we have

$$\frac{1}{x_k} + \Delta'_t = \beta_{k-1}(x'_1, \dots, x'_{k-2}, x_k).$$

Thus,

$$\beta_k(x_1, \dots, x_k) = \beta_{k-1}(x'_1, \dots, x'_{k-2}, x_k) + \Delta_t. \quad (9)$$

Since S is gcd-closed, $(x_k, x_t) \in S$. Let $x_l = (x_k, x_t)$. Assume that $x_l = x_t$. Then $x_t \mid x_k$. Thus $x_t \notin T_k$. It is a contradiction. So $x_l < x_t$, i.e., $l < t$. Therefore,

$$\begin{aligned}
 \Delta_t &= \sum_{r=0}^{k-3} (-1)^{r+1} \cdot \sum_{\substack{1 \leq i_1 < \dots < i_r \leq k-1 \\ i_j \neq t, \\ i_j \neq l.}} \frac{1}{(x_{i_1}, \dots, x_{i_r}, x_t, x_k)} + \sum_{r=0}^{k-3} (-1)^{r+2} \\
 &\quad \cdot \sum_{\substack{1 \leq i_1 < \dots < i_r \leq k-1 \\ i_j \neq t, \\ i_j \neq l.}} \frac{1}{(x_{i_1}, \dots, x_{i_r}, x_l, x_t, x_k)} \\
 &= \sum_{r=0}^{k-3} \sum_{\substack{1 \leq i_1 < \dots < i_r \leq k-1 \\ i_j \neq t, \\ i_j \neq l.}} \left(\frac{(-1)^{r+1}}{(x_{i_1}, \dots, x_{i_r}, x_t, x_k)} \right. \\
 &\quad \left. + \frac{(-1)^{r+2}}{(x_{i_1}, \dots, x_{i_r}, x_l, x_t, x_k)} \right) \\
 &= \sum_{r=0}^{k-3} \sum_{\substack{1 \leq i_1 < \dots < i_r \leq k-1 \\ i_j \neq t, \\ i_j \neq l.}} \left(\frac{(-1)^{r+1}}{(x_{i_1}, \dots, x_{i_r}, x_l)} + \frac{(-1)^{r+2}}{(x_{i_1}, \dots, x_{i_r}, x_l)} \right) \\
 &= 0.
 \end{aligned}$$

By (9) we have that

$$\beta_k(x_1, \dots, x_k) = \beta_{k-1}(x'_1, \dots, x'_{k-2}, x_k), \quad (10)$$

where $\{x'_1, \dots, x'_{k-2}, x_k\} = \{x_1, \dots, x_k\} \setminus \{x_t\}$.

If $T_k \setminus \{x_t\} = \phi$, i.e., $T_k = \{x_t\}$, then it follows that (5) holds from (10) and (3). If $T_k \setminus \{x_t\} \neq \phi$, then the above reduction can continue. Note that T_k is finite. So we can repeat $|T_k|$ times the above reduction. This completes the proof. ■

DEFINITION. Let T be a set of distinct positive integers. For any $a, b \in T$ and $a < b$, we say that a is the greatest-type divisor of b in T , if $a \mid b$ and it can be deduced that $c = a$ from $a \mid c$, $c \mid b$, $c < b$, and $c \in T$.

For example, $T = \{1, 2, 3, 4, 5, 6, 8\}$. Then 2 is the greatest-type divisor of 4 and 6 in T , but 2 is not the greatest-type divisor of 8 in T .

LEMMA 3. Let $R_k = \{y_{k,1}, \dots, y_{k,l_k}\}$ be the set of the greatest-type divisors of x_k in S , where $y_{k,1} < \dots < y_{k,l_k}$, $l_1 = 0$, $l_2 = l_3 = 1$, and $1 \leq l_k \leq k - 2$ for $k \geq 4$. Then

$$\alpha_k = \beta_{l_k+1}(y_{k,1}, \dots, y_{k,l_k}, x_k). \quad (11)$$

Proof. If $k = 1$, then $R_1 = \phi$. So $\alpha_1 = \beta_1(x_1)$. Thus (11) holds. If $k = 2$, then $R_1 = \{x_1\}$. By (3) we have that $\alpha_2 = \beta_2(x_1, x_2)$. Then (11) holds. In the following we let $k \geq 3$. Let $T_k = \{x \in S \mid x < x_k, x \nmid x_k\}$, $S'_k = \{x_1, \dots, x_k\} \setminus T_k$. Write $S'_k = \{x'_1, \dots, x'_h\}$, where $x_1 = x'_1 < \dots < x'_h = x_k$ and $2 \leq h = k - |T_k| \leq k$. Then $R_k \subset S'_k$ and R_k is equal to the set of the greatest-type divisors of x_k in S'_k . If $h = 2$, then $S'_k = \{x_1, x_k\}$ and $T_k = \{x_2, \dots, x_{k-1}\}$. Thus $R_k = \{x_1\}$. By (5) we have that $\alpha_k = \beta_2(x_1, x_k)$ so (11) holds. In the following we let $h \geq 3$. We claim that $\{x'_1, \dots, x'_{h-1}\} \setminus R_k \neq \phi$. Otherwise, $R_k = \{x'_1, \dots, x'_{h-1}\}$. Since $x_1 | x_i$ ($i > 1$) (since S is gcd-closed), one has $x'_1 | x_i$ ($i > 1$). Note that $x'_{h-1} \in \{x_2, \dots, x_{k-1}\}$. So $x'_1 | x'_{h-1}$. Then $x'_1 \notin R_k$. It is a contradiction. Thus the claim is true.

Let $x'_i \in \{x'_1, \dots, x'_{h-1}\} \setminus R_k$. Then $x'_i \notin R_k$. So there exists $x \in \{x'_1, \dots, x'_{h-1}\} \setminus \{x'_i\}$, such that $x'_i | x$. Let $R = \{z_1, \dots, z_p\}$ ($p \geq 1$) be the set of such element x . Namely, $\{z_1, \dots, z_p\} = \{x \mid x \in \{x'_1, \dots, x'_{h-1}\} \setminus \{x'_i\}, \text{ and } x'_i | x\}$. Then $x'_i | z_i$ ($1 \leq i \leq p$). Let $\{w_1, \dots, w_q\} = \{x'_1, \dots, x'_{h-1}\} \setminus (R \cup \{x'_i\})$, where $q = h - p - 2$. By Lemma 2 we have that

$$\alpha_k = \beta_h(x'_1, \dots, x'_h).$$

Note that $x'_h = x_k$. In a way similar to that in (6), we have that

$$\alpha_k = \frac{1}{x_k} + \overline{\Delta}_t + \overline{\Delta}_t, \quad (12)$$

where

$$\overline{\Delta}_t = \sum_{r=1}^{h-2} (-1)^r \cdot \sum_{\substack{1 \leq i_1 < \dots < i_r \leq h-1 \\ i_j \neq t}} \frac{1}{(x'_{i_1}, \dots, x'_{i_r}, x_k)}, \quad (13)$$

$$\overline{\Delta}_t = \sum_{r=0}^{h-2} (-1)^{r+1} \cdot \sum_{\substack{1 \leq i_1 < \dots < i_r \leq h-1 \\ i_j \neq t}} \frac{1}{(x'_{i_1}, \dots, x'_{i_r}, x'_t, x_k)}. \quad (14)$$

For $\overline{\Delta}_t$ we have

$$\begin{aligned} \overline{\Delta}_t &= \sum_{r=0}^q (-1)^{r+1} \sum_{1 \leq i_1 < \dots < i_r \leq q} \frac{1}{(w_{i_1}, \dots, w_{i_r}, x'_t, x_k)} \\ &+ \sum_{r=0}^q \sum_{e=1}^p (-1)^{r+e+1} \cdot \sum_{\substack{1 \leq i_1 < \dots < i_r \leq q, \\ 1 \leq j_1 < \dots < j_e \leq p}} \frac{1}{(w_{i_1}, \dots, w_{i_r}, z_{j_1}, \dots, z_{j_e}, x'_t, x_k)}. \end{aligned} \quad (15)$$

Since $x'_t | z_i$ ($1 \leq i \leq p$), we have that $(z_{j_1}, \dots, z_{j_e}, x'_t) = x'_t$ for any $1 \leq j_1 < \dots < j_e \leq p$. So we have that $(w_{i_1}, \dots, w_{i_r}, x'_t, x_k) = (w_{i_1}, \dots, w_{i_r}, z_{j_1}, \dots, z_{j_e}, x'_t, x_k)$ for any $1 \leq j_1 < \dots < j_e \leq p$ and $1 \leq i_1 < \dots < i_r \leq q$. Then, by (15), we have that

$$\begin{aligned} \overline{\Delta}_t &= \sum_{r=0}^q (-1)^{r+1} \sum_{1 \leq i_1 < \dots < i_r \leq q} \frac{1}{(w_{i_1}, \dots, w_{i_r}, x'_t, x_k)} \\ &+ \sum_{r=0}^q \sum_{e=1}^p (-1)^{r+e+1} \cdot \sum_{\substack{1 \leq i_1 < \dots < i_r \leq q, \\ 1 \leq j_1 < \dots < j_e \leq p}} \frac{1}{(w_{i_1}, \dots, w_{i_r}, x'_t, x_k)} \\ &= \sum_{r=0}^q (-1)^{r+1} \cdot \sum_{1 \leq i_1 < \dots < i_r \leq q} \frac{1}{(w_{i_1}, \dots, w_{i_r}, x'_t, x_k)} \\ &\quad \times \left(1 + \sum_{e=1}^p (-1)^e \cdot \sum_{1 \leq j_1 < \dots < j_e \leq p} 1 \right) \\ &= \sum_{r=0}^q (-1)^{r+1} \cdot \sum_{1 \leq i_1 < \dots < i_r \leq q} \frac{1}{(w_{i_1}, \dots, w_{i_r}, x'_t, x_k)} \\ &\quad \times \left(1 + \sum_{e=1}^p (-1)^e \cdot \binom{p}{e} \right) \\ &= \sum_{r=0}^q (-1)^{r+1} \cdot \sum_{1 \leq i_1 < \dots < i_r \leq q} \frac{1}{(w_{i_1}, \dots, w_{i_r}, x'_t, x_k)} \cdot (1-1)^p \\ &= 0. \end{aligned}$$

By (12), and the definition of β_{h-1} , we have that

$$\alpha_k = \beta_{h-1}(x'_1, \dots, x'_{t-1}, x'_{t+1}, \dots, x'_h). \quad (16)$$

If $\{x'_1, \dots, x'_{h-1}\} \setminus (R_k \cup \{x'_t\}) = \emptyset$, then $R_k = \{x'_1, \dots, x'_{h-1}\} \setminus \{x'_t\}$. So (11) holds by (16). If $\{x'_1, \dots, x'_{h-1}\} \setminus (R_k \cup \{x'_t\}) \neq \emptyset$, then the above

reduction can continue. Let $L_k = \{x'_1, \dots, x'_{h-1}\} \setminus R_k$. Then L_k is finite. So (11) holds by using $|L_k|$ times the above reduction. This completes the proof. ■

In the following lemmas let $R_k = \{y_1, y_2, \dots, y_m\}$ be the set of the greatest-type divisors of x_k ($1 \leq k \leq n$) in S , where $y_1 < y_2 < \dots < y_m$. Then $R_1 = \phi$ and $R_k \neq \phi$ for $k \geq 2$. If $m \geq 2$, we suppose that $G = (y_1, \dots, y_m)$ and $y_i = G \cdot y'_i$ ($1 \leq i \leq m$). Then $1 < y'_1 < y'_2 < \dots < y'_m$ and $(y'_1, y'_2, \dots, y'_m) = 1$. Define $M^{(m)} = \bigcup_{r=2}^m M_r^{(m)}$, where $M_r^{(m)} = \{(y_{i_1}, \dots, y_{i_r}) | 1 \leq i_1 < \dots < i_r \leq m\}$ ($2 \leq r \leq m$), and the $(y_{i_1}, \dots, y_{i_r})$ denotes the greatest common divisor of y_{i_1}, \dots, y_{i_r} . So $G \in M^{(m)}$ and $|M^{(m)}| \geq 1$. Thus $G|x$ for any $x \in M^{(m)}$.

LEMMA 4. Let positive integer $m \leq 2$. Then $\beta_{m+1}(y_1, \dots, y_m, x_k) \neq 0$.

Proof. If $m = 1$, then $\beta_2(y_1, x_k) = (1/x_k) - (1/y_1)$ by (4). Since $y_1 < x_k$, $\beta_2(y_1, x_k) < 0$. If $m = 2$, noting that $(y_1, y_2) = G$, then by (4) we have that

$$\begin{aligned} \beta_3(y_1, y_2, x_k) &= \frac{1}{x_k} - \frac{1}{y_1} - \frac{1}{y_2} + \frac{1}{G} \\ &= \frac{1}{x_k} + \frac{1}{G} \left(1 - \frac{1}{y'_1} - \frac{1}{y'_2} \right). \end{aligned} \quad (17)$$

Since $y'_1 \geq 2$ and $y'_2 \geq 3$,

$$1 - \frac{1}{y'_1} - \frac{1}{y'_2} \geq 1 - \frac{1}{2} - \frac{1}{3} > 0.$$

Thus by (17) we have that $\beta_3(y_1, y_2, x_k) > 0$. The proof is complete. ■

LEMMA 5. Let $m \geq 3$. If $|M^{(m)}| = 1$, then $\beta_{m+1}(y_1, \dots, y_m, x_k) \neq 0$.

Proof. Since $G \in M^{(m)}$ and $|M^{(m)}| = 1$, $M^{(m)} = \{G\}$. Then we have that $(y_{i_1}, \dots, y_{i_r}) = G$ for any $1 \leq i_1 < \dots < i_r \leq m$ ($2 \leq r \leq m$). It implies that $(y'_{i_1}, \dots, y'_{i_r}) = 1$. By (4) we have that

$$\begin{aligned} \beta_{m+1}(y_1, \dots, y_m, x_k) &= \frac{1}{x_k} - \sum_{i=1}^m \frac{1}{y_i} + \sum_{r=2}^m (-1)^r \cdot \sum_{1 \leq i_1 < \dots < i_r \leq m} \frac{1}{G} \\ &= \frac{1}{x_k} - \sum_{i=1}^m \frac{1}{y_i} + \frac{1}{G} \cdot \sum_{r=2}^m (-1)^r \cdot \binom{m}{r} \\ &= \frac{1}{x_k} + \frac{1}{G} \left(m - 1 - \sum_{i=1}^m \frac{1}{y'_i} \right). \end{aligned} \quad (18)$$

Since $y'_2 > y'_1 \geq 2$,

$$1 - \frac{1}{y'_1} - \frac{1}{y'_2} > 0.$$

For $i = 3, \dots, m$, we have that $1 - 1/y'_i > 0$. So

$$m - 1 - \sum_{i=1}^m \frac{1}{y'_i} = \left(1 - \frac{1}{y'_1} - \frac{1}{y'_2}\right) + \sum_{i=3}^m \left(1 - \frac{1}{y'_i}\right) > 0.$$

By (18) we have that $B_{m+1}(y_1, \dots, y_m, x_k) > 0$. This completes the proof. \blacksquare

LEMMA 6. *If $|M^{(3)}| \leq 3$, then $\beta_4(y_1, y_2, y_3, x_k) \neq 0$.*

Proof. By Lemma 5, we need only to consider the case $|M^{(3)}| = 2$ and the case $|M^{(3)}| = 3$.

If $|M^{(3)}| = 2$, then let $M^{(3)} = \{G, x \cdot G\}$, where $x > 1$. Then at least one of the three elements (y_i, y_j) , $1 \leq i < j \leq 3$, is $x \cdot G$. Since $(y_1, y_2, y_3) = G$, so, exactly, one of the three elements (y_i, y_j) , $1 \leq i < j \leq 3$, is $x \cdot G$, and the other two elements are G . By (4) we have that

$$\begin{aligned} \beta_4(y_1, y_2, y_3, x_k) &= \frac{1}{x_k} - \sum_{i=1}^3 \frac{1}{y_i} + \frac{1}{x \cdot G} + \frac{2}{G} - \frac{1}{G} \\ &= \frac{1}{x_k} - \sum_{i=1}^3 \frac{1}{y_i} + \frac{1}{x \cdot G} + \frac{1}{G} \\ &= \frac{1}{x_k} + \frac{1}{G} \left(- \sum_{i=1}^3 \frac{1}{y'_i} + \frac{1}{x} + 1 \right). \end{aligned} \quad (19)$$

Let $(y_a, y_b) = x \cdot G$ and $(y_a, y_c) = (y_b, y_c) = G$, where $1 \leq a, b, c \leq 3$, and a, b , and c are distinct. Since y_a, y_b are the greatest-type divisors and $(y'_a, y'_b) = x$,

$$-\frac{1}{y'_a} - \frac{1}{y'_b} + \frac{1}{x} \geq -\frac{1}{2x} - \frac{1}{3x} + \frac{1}{x} > 0.$$

Note that $-1/y'_c + 1 > 0$. Then

$$- \sum_{i=1}^3 \frac{1}{y'_i} + \frac{1}{x} + 1 = \left(-\frac{1}{y'_a} - \frac{1}{y'_b} + \frac{1}{x} \right) + \left(-\frac{1}{y'_c} + 1 \right) > 0.$$

By (19) we have that $\beta_4(y_1, y_2, y_3, x_k) > 0$.

If $|M^{(3)}| = 3$, then let $M^{(3)} = \{G, x \cdot G, y \cdot G\}$, where $1 < x < y$. Then there are two of the three elements (y_i, y_j) , $1 \leq i < j \leq 3$, such that one is $x \cdot G$ and the other is $y \cdot G$. It follows from $(y_1, y_2, y_3) = G$ that $(x, y) = 1$. Furthermore, exactly one of the three elements (y_i, y_j) , $1 \leq i < j \leq 3$, is $x \cdot G$, one is $y \cdot G$, and one is G . By (4) we have that

$$\begin{aligned}\beta_4(y_1, y_2, y_3, x_k) &= \frac{1}{x_k} - \sum_{i=1}^3 \frac{1}{y_i} + \frac{1}{x \cdot G} + \frac{1}{y \cdot G} + \frac{1}{G} - \frac{1}{G} \\ &= \frac{1}{x_k} - \sum_{i=1}^3 \frac{1}{y_i} + \frac{1}{x \cdot G} + \frac{1}{y \cdot G} \\ &= \frac{1}{x_k} + \frac{1}{G} \cdot \left(- \sum_{i=1}^3 \frac{1}{y'_i} + \frac{1}{x} + \frac{1}{y} \right).\end{aligned}\quad (20)$$

Let $(y_d, y_e) = x \cdot G$, $(y_d, y_f) = y \cdot G$, where $1 \leq d, e, f \leq 3$ and d, e, f are distinct. Since y_d, y_e are the greatest-type divisors and $(y'_d, y'_e) = x$,

$$-\frac{1}{y'_d} - \frac{1}{y'_e} + \frac{1}{x} \geq -\frac{1}{2x} - \frac{1}{3x} + \frac{1}{x} > 0.$$

Since y_f is the greatest-type divisor and $(y'_d, y'_f) = y$, $y'_f > y$. Then $-1/y'_f + 1/y > 0$. Hence

$$-\sum_{i=1}^3 \frac{1}{y'_i} + \frac{1}{x} + \frac{1}{y} = \left(-\frac{1}{y'_d} - \frac{1}{y'_e} + \frac{1}{x} \right) + \left(-\frac{1}{y'_f} + \frac{1}{y} \right) > 0.$$

By (20) we have $\beta_4(y_1, y_2, y_3, x_k) > 0$. The proof is complete. ■

LEMMA 7. If $|M^{(4)}| \leq 2$, then $\beta_5(y_1, y_2, y_3, y_4, x_k) \neq 0$.

Proof. By Lemma 5, we need only to consider the case $|M^{(4)}| = 2$.

Now let $|M^{(4)}| = 2$. Then $M^{(4)} = \{G, x \cdot G\}$ for some $x > 1$. So at least one of the six elements (y_i, y_j) , $1 \leq i < j \leq 4$, is $x \cdot G$. Since $(y_1, y_2, y_3, y_4) = G$, at most three of the six elements (y_i, y_j) , $1 \leq i < j \leq 4$, are $x \cdot G$, and for $1 \leq i \leq 4$, $1 \leq i_1 < i_2 < i_3 \leq 4$ and $i_j \neq i$ ($j = 1, 2, 3$), we have either $(y_i, y_{i_1}) \neq x \cdot G$, $(y_i, y_{i_2}) \neq x \cdot G$, or $(y_i, y_{i_3}) \neq x \cdot G$. Consider the following cases:

Case 1. If exactly one of the six elements (y_i, y_j) , $1 \leq i < j \leq 4$, is $x \cdot G$, then the other five elements are G . Hence, the four elements

(y_i, y_j, y_l) , $1 \leq i < j < l \leq 4$, are G . By (4) we have that

$$\begin{aligned}\beta_5(y_1, y_2, y_3, y_4, x_k) &= \frac{1}{x_k} - \sum_{i=1}^4 \frac{1}{y_i} + \frac{1}{x \cdot G} + \frac{5}{G} - \frac{4}{G} + \frac{1}{G} \\ &= \frac{1}{x_k} - \sum_{i=1}^4 \frac{1}{y_i} + \frac{1}{x \cdot G} + \frac{2}{G} \\ &= \frac{1}{x_k} + \frac{1}{G} \cdot \left(- \sum_{i=1}^4 \frac{1}{y'_i} + \frac{1}{x} + 2 \right).\end{aligned}\quad (21)$$

Since $2 \leq y'_1 < y'_2 < y'_3 < y'_4$, $-\sum_{i=1}^4 1/y'_i + 2 \geq 2 - \frac{1}{2} - \frac{1}{3} - \frac{1}{4} - \frac{1}{5} > 0$. By (21) we have that $\beta_5(y_1, y_2, y_3, y_4, x_k) > 0$.

Case 2. If at least two of the six elements (y_i, y_j) , $1 \leq i < j \leq 4$, are $x \cdot G$, then there do not exist four distinct positive integers e, f, g, h , $1 \leq e, f, g, h \leq 4$, such that $(y_e, y_f) = (y_g, y_h) = x \cdot G$. Otherwise we have that $(y_1, y_2, y_3, y_4) = (y_e, y_f, y_g, y_h) = ((y_e, y_f), (y_g, y_h)) = x \cdot G$. It contradicts to $(y_1, y_2, y_3, y_4) = G$. So there are only the following four cases:

- (i) $(y_1, y_2) = (y_1, y_3) = x \cdot G$. Then $(y_2, y_3) = x \cdot G$ and the other three elements (y_i, y_j) are G ;
- (ii) $(y_1, y_2) = (y_1, y_4) = x \cdot G$. Then $(y_2, y_4) = x \cdot G$ and the other three elements (y_i, y_j) are G ;
- (iii) $(y_1, y_3) = (y_1, y_4) = x \cdot G$. Then $(y_3, y_4) = x \cdot G$ and the other three elements (y_i, y_j) are G ;
- (iv) $(y_2, y_3) = (y_2, y_4) = x \cdot G$. Then $(y_3, y_4) = x \cdot G$ and the other three elements (y_i, y_j) are G .

Then it can be deduced that exactly three of the six elements (y_i, y_j) , $1 \leq i < j \leq 4$, are $x \cdot G$, and the other three elements are G . Thus, exactly one of the four elements (y_i, y_j, y_l) , $1 \leq i < j < l \leq 4$, is $x \cdot G$, and the other three elements are G . By (4) we have that

$$\begin{aligned}\beta_5(y_1, y_2, y_3, y_4, x_k) &= \frac{1}{x_k} - \sum_{i=1}^4 \frac{1}{y_i} + \frac{3}{x \cdot G} + \frac{3}{G} - \frac{1}{x \cdot G} - \frac{3}{G} + \frac{1}{G} \\ &= \frac{1}{x_k} - \sum_{i=1}^4 \frac{1}{y_i} + \frac{2}{x \cdot G} + \frac{1}{G} \\ &= \frac{1}{x_k} + \frac{1}{G} \cdot \left(- \sum_{i=1}^4 \frac{1}{y'_i} + \frac{2}{x} + 1 \right).\end{aligned}\quad (22)$$

Let $(y_a, y_b) = (y_a, y_c) = (y_b, y_c) = x \cdot G$, where $1 \leq a, b, c \leq 4$, and a, b, c are distinct. Since y_a, y_b, y_c are the greatest-type divisors of x_k , $\{y'_a, y'_b, y'_c\}$ has one element not less than $2x$, one not less than $3x$, and one not less than $4x$. Thus

$$-\frac{1}{y'_a} - \frac{1}{y'_b} - \frac{1}{y'_c} + \frac{2}{x} \geq \frac{2}{x} - \frac{1}{2x} - \frac{1}{3x} - \frac{1}{4x} > 0.$$

Now let $\{y'_d\} = \{y'_1, y'_2, y'_3, y'_4\} \setminus \{y'_a, y'_b, y'_c\}$. Then $-1/y'_d + 1 > 0$. Therefore,

$$-\sum_{i=1}^4 \frac{1}{y'_i} + \frac{2}{x} + 1 = \left(-\frac{1}{y'_a} - \frac{1}{y'_b} - \frac{1}{y'_c} + \frac{2}{x}\right) + \left(-\frac{1}{y'_d} + 1\right) > 0.$$

By (22) we have that $\beta_5(y_1, y_2, y_3, y_4, x_k) > 0$. The proof is complete. ■

III. THE MAIN THEOREM

In this section, we give the main result of this paper as follows.

THEOREM. *Let n be a positive integer.*

(i) *If $n \leq 7$, then the Bourque–Ligh's conjecture is true. Namely, for any gcd-closed set $S = \{x_1, \dots, x_n\}$ of n distinct positive integers, the LCM matrix $[S]_n$ defined on S is nonsingular.*

(ii) *If $n \geq 8$, then there exists a gcd-closed set $S = \{x_1, \dots, x_n\}$ of n distinct positive integers, such that $\alpha_n(x_1, \dots, x_n) = 0$. Therefore the Bourque–Ligh conjecture is not true for $n \geq 8$.*

Proof. (i) Let $S = \{x_1, \dots, x_n\}$ be a gcd-closed set of n distinct positive integers. Without loss of generality, we may let $1 \leq x_1 < x_2 < \dots < x_n$. For $1 \leq k \leq n$, let $R_k = \{y_1, \dots, y_m\}$ be the set of the greatest-type divisors of x_k in S , where $y_1 < \dots < y_m$. Then $R_1 = \phi$ and $R_k \neq \phi$ (i.e., $m \geq 1$) for $k \geq 2$. By Lemma 3 we have that

$$\alpha_k = \beta_{m+1}(y_1, \dots, y_m, x_k). \quad (23)$$

For $m \geq 2$, let $M_r^{(m)} = \{(y_{i_1}, \dots, y_{i_r}) | 1 \leq i_1 < \dots < i_r \leq m\}$ ($2 \leq r \leq m$). Let $M^{(m)} = \bigcup_{r=2}^m M_r^{(m)}$. Clearly, $R_k \subset \{x_1, \dots, x_{k-1}\}$. It follows from S is gcd-closed that $M^{(m)} \subset \{x_1, \dots, x_{k-1}\}$. Since y_1, \dots, y_m are the greatest-type divisors of x_k in S , $R_k \cap M^{(m)} = \phi$. So $m + |M^{(m)}| \leq k - 1$. Then

for $m \geq 2$, we have that

$$1 \leq |M^{(m)}| \leq k - m - 1. \quad (24)$$

We claim that $\alpha_k \neq 0$ for $1 \leq k \leq 7$.

Let $k = 1$. Then $\alpha_1 = 1/x_1 \neq 0$.

Let $k = 2$ or 3 . Then $m = 1$. It follows from (23) and Lemma 4 that $\alpha_k = \beta_2(y_1, x_k) \neq 0$.

Let $k = 4$. Suppose that $m \geq 3$, then by (24) we have that $k \geq m + 2 \geq 3 + 2 = 5$. It is a contradiction and so $m \leq 2$. It follows from (23) and Lemma 4 that $\alpha_4 = \beta_{m+1}(y_1, \dots, y_m, x_4) \neq 0$.

Let $k = 5$. If $m \leq 2$, then it follows from (23) and Lemma 4 that $\alpha_5 = \beta_{m+1}(y_1, \dots, y_m, x_5) \neq 0$. If $m \geq 3$, by (24) we have that $|M^{(m)}| \leq 5 - m - 1 \leq 1$. It can be deduced that $|M^{(m)}| = 1$ and $m = 3$. It follows from (23) and Lemma 5 that $\alpha_5 = \beta_4(y_1, y_2, y_3, x_5) \neq 0$.

Let $k = 6$. If $m \leq 2$, then it follows from (23) and Lemma 4 that $\alpha_6 = \beta_{m+1}(y_1, \dots, y_m, x_6) \neq 0$. If $m \geq 3$, by (24) we have that $m \leq k - 2 = 4$. Then $m = 3$ or 4 . If $m = 3$, by (24) we have that $|M^{(3)}| \leq 6 - 3 - 1 = 2$. It follows from (23) and Lemma 6 that $\alpha_6 = \beta_4(y_1, y_2, y_3, x_6) \neq 0$. If $m = 4$, by (24) we have that $|M^{(4)}| \leq 6 - 4 - 1 = 1$. So $|M^{(4)}| = 1$. It follows from (23) and Lemma 5 that $\alpha_6 = \beta_5(y_1, y_2, y_3, y_4, x_6) \neq 0$.

Let $k = 7$. If $m \leq 2$, then it follows from (23) and Lemma 4 that $\alpha_7 = \beta_{m+1}(y_1, \dots, y_m, x_7) \neq 0$. If $m \geq 3$, by (24) we have that $m \leq k - 2 = 5$. Then $m = 3, 4$, or 5 . If $m = 3$, by (24) we have that $|M^{(3)}| \leq 7 - 3 - 1 = 3$. It follows from (23) and Lemma 6 that $\alpha_7 = \beta_4(y_1, y_2, y_3, x_7) \neq 0$. If $m = 4$, by (24) we have that $|M^{(4)}| \leq 7 - 4 - 1 = 2$. Then it follows from (23) and Lemma 7 that $\alpha_7 = \beta_5(y_1, y_2, y_3, y_4, x_7) \neq 0$. If $m = 5$, by (24) we have that $|M^{(5)}| \leq 1$. So $|M^{(5)}| = 1$. Then it follows from (23) and Lemma 5 that $\alpha_7 = \beta_6(y_1, y_2, y_3, y_4, y_5, x_7) \neq 0$.

Therefore the claim is true.

Now let $n \leq 7$. Then for k , $1 \leq k \leq n$, it follows from the claim that $\alpha_k \neq 0$. So by (1) we have that $\det[S]_n \neq 0$. Hence the LCM matrix $[S]_n$ defined on S is nonsingular. Thus the Bourque-Ligh conjecture is true for $n \leq 7$.

(ii) Let $n \geq 8$ and let $a > 1$ be any integer. Now let

$$\begin{aligned} x_i &= a^{i-1}, & 1 \leq i \leq n-7, \\ x_{n-6} &= 2b, & x_{n-5} = 3b, & x_{n-4} = 5b, & x_{n-3} = 36b, \\ x_{n-2} &= 230b, & x_{n-1} = 825b, & x_n = 227700b, \end{aligned}$$

where $b = a^{n-8}$. Then $S = \{x_1, \dots, x_n\}$ be a gcd-closed set of n distinct positive integers. It is easy to see that $\{x_{n-3}, x_{n-2}, x_{n-1}\}$ is the set of the

greatest-type divisors of $x_n = 227700b$ in S . Since

$$\begin{aligned} \beta_4(x_{n-3}, x_{n-2}, x_{n-1}, x_n) &= \frac{1}{b} \cdot \beta_4(36, 230, 825, 227700) \\ &= \frac{1}{b} \cdot \left(\frac{1}{227700} - \frac{1}{36} - \frac{1}{230} - \frac{1}{825} + \frac{1}{2} + \frac{1}{3} + \frac{1}{5} - 1 \right) \\ &= 0. \end{aligned}$$

It follows from Lemma 3 that $\alpha_n = \beta_4(x_{n-3}, x_{n-2}, x_{n-1}, x_n) = 0$. By (1) we have that $\det[S]_n = 0$, so the LCM matrix $[S]_n$ defined on S is singular. Thus the Bourque–Ligh conjecture is not true for $n \geq 8$. The proof is complete. ■

ACKNOWLEDGMENT

The author is very grateful to Professor Sun Qi for his help and encouragement in the writing of this paper.

REFERENCES

1. T. M. Apostol, Arithmetical properties of generalized Ramanujan sums, *Pacific J. Math.* **41** (1972), 281–293.
2. S. Beslin and S. Ligh, Greatest common divisor matrices, *Linear Algebra Appl.* **118** (1989), 69–76.
3. K. Bourque and S. Ligh, On GCD and LCM matrices, *Linear Algebra Appl.* **174** (1992), 65–74.
4. K. Bourque and S. Ligh, Matrices associated with classes of arithmetical functions, *J. Number Theory* **45** (1993), 367–376.
5. P. Haukkanen, J. Wang, and J. Sillanpää, On Smith's determinant, submitted for publication.
6. S. Hong, LCM matrix on an r -fold gcd-closed set, *J. Sichuan University (Natu. Sci. Ed.)* **33**, No. 6 (1996), 650–657.
7. P. J. McCarthy, A generalization of Smith's determinant, *Canad. Math. Bull.* **29** (1988), 109–113.
8. H. J. S. Smith, On the value of a certain arithmetical determinant, *Proc. London Math. Soc.* **7** (1875–1876), 208–212.